

The Nullstellensatz for Sparse Tropical Polynomial Systems

Antoine Béreau (antoine.bereau@inria.fr)

supervised by Marianne Akian and Stéphane Gaubert

CMAP (École polytechnique), IP Paris, CNRS, Inria

We fix a collection $f = (f_1, \dots, f_k)$ of tropical polynomials in n variables with respective degrees (d_1, \dots, d_k) .

Motivation

Theorem (Tropical Dual Nullstellensatz, [GP18]) The polynomials of f have a common root $x \in \mathbb{R}^n$ iff there exists a vector $y \in \mathbb{R}^m$ with $m = \binom{N+n}{n}$ in the tropical right null space of the truncated Macaulay matrix \mathcal{M}_N for

$$N = (n+2)(d_1 + \dots + d_k).$$

Question: What is the smallest possible value of N such that this result holds and how can this result be expressed for **sparse** polynomials?

Tropical Polynomials

• **Tropical semiring:** $\mathbb{R}_\infty = (\mathbb{R} \cup \{-\infty\}, \oplus, \odot)$ with $\oplus := \max, \odot := +$, zero element $\mathbb{0} := -\infty$ and unit element $\mathbb{1} := 0$.

• $x \in \mathbb{R}_\infty^n$ is a **root** of a tropical (Laurent) polynomial p whenever the maximum in the expression

$$p(x) = \bigoplus_{\alpha \in A} p_\alpha \odot x^{\odot \alpha} = \max_{\alpha \in A} (p_\alpha + \langle x, \alpha \rangle)$$

is attained twice, where $A \subset \mathbb{Z}^n$ is the support of p . This is denoted as $p(x) \nabla \mathbb{0}$.

• y is in the **tropical right null space** of a $\ell \times m$ matrix $A = (a_{ij})$ whenever for all $1 \leq i \leq \ell$, the maximum in the expression

$$\bigoplus_{j=1}^m a_{ij} \odot y_j = \max_{1 \leq j \leq m} (a_{ij} + y_j)$$

is attained twice. This is denoted as $A \odot y \nabla \mathbb{0}$.

The Macaulay matrix

• The **Macaulay matrix** associated to f is the (infinite) matrix $\mathcal{M} = (m_{(i,\alpha),\beta})$ indexed by $([n] \times \mathbb{Z}^n) \times \mathbb{Z}^n$, where $m_{(i,\alpha),\beta}$ corresponds to the coefficient of X^β in the tropical polynomial $X^\alpha f_i$.

• A finite subset \mathcal{E} of \mathbb{Z}^n yields a (finite) submatrix $\mathcal{M}_\mathcal{E}$ of \mathcal{M} obtained by taking only the rows whose support is included in \mathcal{E} and the columns indexed by \mathcal{E} .

• For $\mathcal{E} = \{\alpha \in \mathbb{N}^n : \alpha_1 + \dots + \alpha_n \leq N\}$, we denote $\mathcal{M}_N := \mathcal{M}_\mathcal{E}$.

Newton polytopes

• Set for $1 \leq i \leq k$, $Q_i := \text{conv}(\mathcal{A}_i)$ the **Newton polytope** of f_i and $Q = Q_1 + \dots + Q_k$.

• The upper hull of the lifted support $\{(\alpha, f_{i,\alpha}) : \alpha \in \mathcal{A}_i\}$ is the graph of a function h_i with support Q_i and if $h := h_1 \square \dots \square h_k$ where \square denotes the sup-convolution, then $\text{hypo}(h) = \text{hypo}(h_1) + \dots + \text{hypo}(h_k)$.

• The projection of $\text{hypo}(h)$ onto Q yields a coherent mixed subdivision of Q .


Canny-Emiris subsets of \mathbb{Z}^n

Canny-Emiris set associated to f : any set of the form $\mathcal{E} = (Q + \delta) \cap \mathbb{Z}^n$ with δ a generic vector in the linear space directing the affine hull of Q .

Examples

$$\text{Consider } (E_1) : \begin{cases} f_1 = 1 \oplus 2x_1 \oplus 1x_2 \oplus 1x_1x_2 \\ f_2 = 0 \oplus 0x_1 \oplus 1x_2 \\ f_3 = 2x_1 \oplus 0x_2 \end{cases} \text{ and } (E_2) : \begin{cases} f_1 = 1 \oplus 4x_1 \oplus 1x_2 \oplus 3x_1x_2 \\ f_2 = 0 \oplus 0x_1 \oplus 1x_2 \\ f_3 = 2x_1 \oplus 0x_2 \end{cases}$$

The Newton polytopes associated to both systems and their Minkowski sum are as follow.




For $\delta = (-1 + \varepsilon, -1 + \varepsilon)$ with $\varepsilon > 0$ sufficiently small, we obtain the Canny-Emiris set

$$\mathcal{E} := (Q + \delta) \cap \mathbb{Z}^n = \{(0,0), (1,0), (0,1), (2,0), (1,1), (0,2)\}$$


corresponding to the set of monomials $\{1, x_1, x_2, x_1^2, x_1x_2, x_2^2\}$. The associated matrices are

$$\mathcal{M}_\mathcal{E}^{(1)} = \begin{matrix} & 1 & x_1 & x_2 & x_1^2 & x_1x_2 & x_2^2 \\ \begin{matrix} f_1 \\ f_2 \\ f_3 \end{matrix} & \begin{pmatrix} 1 & 2 & 1 & 1 \\ 0 & 0 & 1 \\ 2 & 0 & 0 & 1 \end{pmatrix} \end{matrix} \text{ and } \mathcal{M}_\mathcal{E}^{(2)} = \begin{matrix} & 1 & x_1 & x_2 & x_1^2 & x_1x_2 & x_2^2 \\ \begin{matrix} f_1 \\ f_2 \\ f_3 \end{matrix} & \begin{pmatrix} 1 & 4 & 1 & 3 \\ 0 & 0 & 1 \\ 2 & 0 & 0 & 1 \end{pmatrix} \end{matrix}$$


(a) The arrangement of tropical varieties of the polynomials from the system (E_1) .




(b) The subdivision of Q associated to (E_1) .




(c) The arrangement of tropical varieties of the polynomials from the system (E_2) .




(d) The subdivision of Q associated to (E_2) .



(e) The subdivision of Q associated to (E_1) arises from the projection of the Minkowski sum of the hypographs of the h_i .



(f) The polytope $Q + \delta$, with interior integer points labelled by the row content of the cell they belong to for system (E_1) .



$$\text{For the system } (E_1), \text{ we obtain the matrix } \mathcal{M}_\mathcal{E}^{(1)} = \begin{matrix} & (0,0) & (1,0) & (0,1) & (2,0) & (1,1) & (0,2) \\ \begin{matrix} f_1 \\ f_2 \\ f_3 \end{matrix} & \begin{pmatrix} 1 & 2 & 1 & 1 \\ 2 & 0 & 0 & 1 \\ 0 & 0 & 1 \\ 2 & 0 & 0 & 1 \end{pmatrix} \end{matrix} \text{ which is nonsingular.}$$

For the system (E_2) , for $y = \text{ver}(-3, -1) = (0, -3, -1, -6, -4, -2)$, we obtain $\mathcal{M}_\mathcal{E}^{(2)} \odot y \nabla \mathbb{0}$.

Nullstellensatz for Sparse Tropical Polynomial Systems

Theorem The system $f \nabla \mathbb{0}$ has a solution $x \in \mathbb{R}^n$ iff there exists a vector $y \in \mathbb{R}^m$ in the tropical right null space of the submatrix $\mathcal{M}_{\mathcal{E}'}$ of \mathcal{M} , where \mathcal{E}' is any subset of \mathbb{Z}^n containing a nonempty Canny-Emiris set \mathcal{E} .

Corollary The system $f \nabla \mathbb{0}$ has a solution $x \in \mathbb{R}^n$ if and only if the truncated Macaulay tropical linear system $\mathcal{M}_N \odot y \nabla \mathbb{0}$ has a solution $y \in \mathbb{R}^m$ for $N = d_1 + \dots + d_k$.

The Canny-Emiris construction

- If $f \nabla \mathbb{0}$ has a solution $x \in \mathbb{R}^n$, then $y = (x_p)_{p \in \mathcal{E}'}$ of x is a solution to $\mathcal{M}_{\mathcal{E}'} \odot y \nabla \mathbb{0}$.
- Otherwise we apply the Canny-Emiris construction from [CE93] and [Stu94] but in a potentially **non generic** case. If $p \in Q$, then $(p - \delta, h(p - \delta))$ is in the relative interior of a facet F of $\text{hypo}(h)$, and F can be written as $F_1 + \dots + F_k$ with F_i faces of $\text{hypo}(h_i)$.
- Since f does not have a common root, at least one F_i is a singleton. Consider the maximal index j such that $F_j = \{a_j\}$ is a singleton. The couple (j, a_j) is called the **row content** of p .
- If $p \in \mathcal{E}$ and if (j, a_j) is its row content, then the support of the polynomial $X^{p-a_j} f_j$ is included in \mathcal{E} . This allows us to construct a square submatrix $\mathcal{M}_{\mathcal{E}\mathcal{E}} = (m_{pp'})_{(p,p') \in \mathcal{E} \times \mathcal{E}}$ of $\mathcal{M}_\mathcal{E}$.
- The matrix $\tilde{\mathcal{M}}_{\mathcal{E}\mathcal{E}} = (\tilde{m}_{pp'})_{(p,p') \in \mathcal{E} \times \mathcal{E}}$ obtained by setting $\tilde{m}_{pp'} = m_{pp'} - h(p' - \delta)$ is **tropically diagonally dominant**, and therefore its tropical right null space is reduced to $\{\mathbb{0}\}$, and thus this is also the case for $\mathcal{M}_{\mathcal{E}\mathcal{E}}$, hence there does not exist $y \in \mathbb{R}^{\mathcal{E}}$ such that $\mathcal{M}_\mathcal{E} \odot y \nabla \mathbb{0}$.

Perspectives and related results

- The optimal value of N seems to be $d_1 + \dots + d_k - \max(n, k - 1)$.
- **Tropical resultant polynomial** based on [JY13] and generalizing tropical Cramer's theorem from [AGG08] in the case $k = n + 1$.
- **Positivstellensatz** for systems of the form $f^+ = f^-$ and $f^+ \geq f^-$.
- Generalization to any tropical semiring obtained from a totally ordered group with bottom element.

References

- [AGG08] Marianne Akian, Stéphane Gaubert and Alexander Guterman, Linear independence over tropical semirings and beyond, *Contemp. Math.* (2008) **495**
- [CE93] John Canny and Ioannis Emiris, An efficient algorithm for the sparse mixed resultant (1993) **263**:89-104
- [GP18] Dima Grigoriev and Vladimir V. Podolskii, Tropical Effective Primary and Dual Nullstellensätze, *Discrete Comput Geom* (2018) **59**:507-522
- [JY13] Anders Jensen and Josephine Yu, Computing tropical resultants, *Journal of Algebra* (2013) **387**:287-319.
- [Stu94] Bernd Sturmfels, On the Newton Polytope of the Resultant, *Journal of Algebraic Combinatorics* (1994) **3**(2):207-236

